

# Computation of Mass Outflow Rate from Relativistic Quasi-Spherical Accretion onto Black Holes

Tapas K. Das

*Theoretical Astrophysics Group*

*S. N. Bose National Centre For Basic Sciences*

*Block JD Sector III Salt Lake Calcutta 700 091 India*

*E-mail tdas@boson.bose.res.in*

Accepted 08.04.99 Received 07.12.98. ; in original form 07.12.98

## ABSTRACT

We compute mass outflow rate  $R_{\dot{m}}$  from relativistic matter accreting quasi-spherically onto Schwarzschild black holes. Taking the pair-plasma pressure mediated shock surface as the *effective* boundary layer (of the black hole) from where bulk of the outflow is assumed to be generated, computation of this rate is done using combinations of exact transonic inflow and outflow solutions. We find that  $R_{\dot{m}}$  depends on the initial parameters of the flow, the polytropic index of matter, the degree of compression of matter near the shock surface and on the location of the shock surface itself. We thus not only study the variation of the mass outflow rate as a function of various physical parameters governing the problem but also provide a sufficiently plausible estimation of this rate.

**Key words:** AGN – quasars – jets – accretion, accretion discs – black hole physics – shock-waves – hydrodynamics – outflow – wind

**To appear in MNRAS**

## 1 INTRODUCTION

Computation of mass outflow rates from the advective accretion disks around black holes and neutron stars very recently has been done (Chakrabarti, 1998; Das, 1998, 1998a; Das & Chakrabarti, 1998 [DC98 hereafter]). by self consistently combining the exact transonic accretion and wind solutions. Rigorously justifying the fact that most of the outflowing matter comes out from the CENTrifugal Pressure Supported BOundary Layer (CENBOL) it has been shown there that the Rankine-Hugoniot shock location or the location of the maximum polytropic pressure acts as the CENBOL. However, for some black hole models of active galactic nuclei, inflow may not have accretion disk (Rees, 1977 and references therein). Accretion is then quasi-spherical having almost zero or negligible angular momentum (Bondi [1952] type accretion) and the shock is not of Rankine-Hugoniot type. On the otherhand, absence of angular momentum rules out the possibility of formation of the polytropic pressure maxima. So for quasi-spherical accretion, absence of intrinsic angular momentum of the accreting material does not allow CENBOL formation. It has been shown that (Meszaros

& Ostriker, 1983; Kazanas & Ellison, 1986 [KE86 hereafter]) for quasi -spherical accretion onto black holes, steady state situation may be developed where a standing collisionless shock may form due to the plasma instabilities and for non-linearity introduced by small density perturbation. This is because, after crossing the sonic point the infalling matter (in plasma form) becomes highly supersonic. Any small perturbation and slowing down of the infall velocity will create a piston and produce a shock. A spherically symmetric shock produced in such a way will accelerate a fraction of the inflowing plasma to relativistic energies. The shock accelerated relativistic particles suffer essentially no Compton loss and are assumed to lose energy only through  $p - p$  collision. These relativistic hadrons are not readily captured by the black hole (Protheros & Kazanas, 1983) rather considerable high energy density of these relativistic protons would be maintained to support a standing, collisionless, spherical shock around the black hole (see KE86 and references therein). Thus a self-supported standing shock may be produced *even* for accretion with zero angular momentum. In this work, we take this pair-plasma pressure mediated shock surface as the alternative of the CENBOL which can be treated as the *effective* physical hard surface which, in principle mimics the ordinary stellar surface regarding the mass outflow. The condition necessary for the develop-

ment and maintenance of such a self supported spherical shock is satisfied for the high Mach number solutions (Ellison & Eichler, 1984). Keeping this in the back of our mind, for our present work, we concentrate only on low energy accretion to obtain high shock Mach number. Considering low energy ( $\mathcal{E} \lesssim 0.001$ ) accretion, we assume that particles accreting toward black hole are shock accelerated via first order Fermi acceleration (Axford, et. all, 1977) producing relativistic protons. Those relativistic protons usually scatter several times before being captured by the black hole. These energised particles, in turn, provide sufficiently outward pressure to support a standing, collisionless shock. A fraction of the energy flux of infalling matter is assumed to be converted into radiation at the shock standoff distance through hadronic ( $p-p$ ) collision and mesonic ( $\pi^\pm, \pi^0$ ) decay. Luminosity produced by this fraction is used to obtain the shock location for the present work. Our approach to formulate the expression for shock location is somewhat similar to that of KE86.

At the shock surface, density of the post-shock material shoots up and velocity falls down, infalling matter starts piling up on the shock surface. The post shock relativistic hadronic pressure then gives a kick to the piled up matter the result of which is the ejection of outflow from the shock surface. For this type of inflow, accretion is known to proceed smoothly after a shock transition, since successful subsonic solutions have been constructed (Flammang, 1982) for accretion onto black holes embedded within normal stars with the boundary condition  $u = c$ ; where  $u$  is the infall velocity of matter and  $c$  is the velocity of light in vacuum. The fraction of energy converted, the shock compression ratio  $R_{comp}$  (in the notation of DC98), along with the ratio of post shock relativistic hadronic pressure to infalling ram pressure at a given shock location are obtained from the steady state shock solution of Ellison and Eichler (Ellison & Eichler, 1984;1985). The shock location as a function of the specific energy  $\mathcal{E}$  of the infalling matter and accretion rate is then self consistently obtained using the above mentioned quantities. We then calculate the amount of mass outflow rate  $R_{in}$  from the shock surface using combination of *exact* transonic inflow outflow solutions and study the dependence of  $R_{in}$  on various physical entities governing the inflow-outflow system. We thus quantitatively compute the mass outflow rate *only* from the inflow parameters. In this way we analytically connect the accretion and wind type topologies self-consistently. As there was no such attempt available in the literature which computes the mass loss rate from zero angular momentum quasi-spherical Bondi type accretion, our work, for the *first time* we believe, could shed light on the nature of the outflow from the models of AGNs with no accretion disks.

We consider polytropic inflow. The outflow is also assumed to be polytropic except the fact that  $\gamma_{outflow}$  is assumed to be less than the  $\gamma_{inflow}$  reason of which will be discussed in the next section. As a fraction of the energy of infalling material is converted into radiation, energy flux of the wind is somewhat less than that of the accretion but is kept constant throughout the outflow.

The plan of the paper is as follows :

In the next Section, we describe our model and present the governing equations for the inflow and the outflow along with the solution procedure. In §3, we present the result of

our computation. Finally, in §4, we draw our conclusions. A preliminary report of this work has been presented elsewhere (Das, 1998b).

## 2 MODEL DESCRIPTION, GOVERNING EQUATIONS AND THE SOLUTION PROCEDURE

### 2.1 Inflow model

We assume that a Schwarzschild type black hole of mass  $10M_\odot$  quasi-spherically accretes low energy ( $\mathcal{E} \lesssim 0.001$ ) fluid obeying polytropic equation of state. The density of the fluid is  $\rho(r)$ ,  $r$  being the radial distance measured in the unit of Schwarzschild radius. We also assume that the accretion rate (in the unit of Eddington rate) with which the fluid is being accreted, is not a function of  $r$ . For simplicity of calculation, we choose geometric unit (unit of length =  $\frac{2GM}{c^2}$  & unit of velocity is  $c$ .  $M$  is the mass of the black hole,  $c$  is the velocity of light and  $G$  is the universal gravitational constant. All other relevant physical quantities can be expressed like wise) to measure all the relevant quantities. We ignore the self gravity of the flow and the calculation is being done using Paczyn'ski-Wiita (Paczyn'ski & Wiita, 1980) potential which mimics surrounding of the Schwarzschild black hole. The equations (in dimensionless geometric unit) governing the inflow are :

a) Conservation of specific energy is given by:

$$\mathcal{E} = \frac{u(r)^2}{2} + na(r)^2 - \frac{1}{2(r-1)} \quad (1)$$

$u(r)$  and  $a(r)$  are the radial and polytropic sound velocities respectively.  $a(r) = \left[ \frac{\gamma p(r)}{\rho(r)} \right]^{\frac{1}{2}}$ ;  $p(r)$  being the polytropic pressure. For a polytropic inflow,  $p(r) = K\rho(r)^\gamma$  where  $K$  is a measure of the entropy which remains constant throughout the flow except at the shock location.  $n$  is the polytropic constant of the inflow  $n = (\gamma - 1)^{-1}$ ,  $\gamma$  being the polytropic index.

b) Mass conservation equation is given by,

$$\dot{M}_{in} = \Theta_{in} \rho(r) u(r) r^2 \quad (2)$$

As already mentioned, we assume that a steady, collisionless shock forms at a distance  $r_{sh}$  (measured in the unit of Schwarzschild radius) due to the instabilities in the plasma flow. We also assume that for our model, the effective thickness of the shock  $\Delta_{sh}$  is small enough compared to the shock standoff distance, i, e,

$$\Delta_{sh} \ll r_{sh}$$

Accreting particles with infall velocity  $u(r)$  are then assumed to be shock accelerated via first order Fermi acceleration. Due to this process, relativistic protons will be produced. These relativistic protons, suffer essentially no Compton loss and hence are not readily swallowed by the black hole. Rather they usually scatter several times before being captured by the black hole thus provide sufficient outward pressure necessary to support the shock. These protons, in turn,

produce pions through post-shock inelastic nuclear collisions

$$p + p \longrightarrow p + p + \pi^\pm + \pi^0$$

Pions generated by this process, decay into relativistic electrons, neutrinos & antineutrinos and produces high energy  $\gamma$  rays.

$$\pi^0 \longrightarrow 2\gamma$$

$$\pi^- \longrightarrow e^- + \bar{\nu}_e$$

$$\pi^+ \longrightarrow e^+ + \nu_e$$

These electrons produce the observed non-thermal radiation by Synchrotron and inverse Compton scattering. The overall efficiency of this mechanism depends largely on the shock location. It has been shown (Eichler, 1979) that almost half of the energy flux that goes into relativistic particles is lost due to neutrinos.

From eqs. (2), the density of infalling matter  $\rho(r)$  comes out to be

$$\rho(r) = \frac{\dot{M}_{in}}{\Theta_{in} u(r) r^2} \quad (3)$$

At the shock, density of matter will shoot up and inflow velocity fall down abruptly. If  $(\rho_-, u_-)$  and  $(\rho_+, u_+)$  are the pre and post-shock densities and velocities respectively, then

$$\frac{\rho_+}{\rho_-} = R_{comp} = \frac{u_-}{u_+} \quad (4)$$

Where  $R_{comp}$  is the shock compression ratio (in the notation of DC98). For high shock Mach number solution (which is compatible with our low energy accretion model), the expression for  $R_{comp}$  can be well approximated as

$$R_{comp} = 1.44 M_{sh}^{\frac{3}{4}} \quad (5)$$

Where  $M_{sh}$  is the shock Mach number and eqs. (5) holds good for  $M_{sh} \gtrsim 4.0$  (Ellison & Eichler, 1985). However, the scattering mean free path of the relativistic hadrons produced by this process are assumed to be small enough so that they could encounter the full shock compression ratio while crossing the shock.

The hadronic interaction characteristic time scale  $\tau_{pp}$  may be expressed as \*

$$\tau_{pp} = \frac{1}{\rho_+ \sigma_{pp} c} \quad (6)$$

Where  $\sigma_{pp}$  is the collision cross section for relativistic protons. Using eqs. (4),  $\tau_{pp}$  can also be expressed as

$$\tau_{pp} = \frac{1}{R_{comp} \rho_- \sigma_{pp} c} \quad (6a)$$

\* As only a fraction of the accreting matter is shock energized, the value of  $\rho_+$  used to calculate  $\tau_{pp}$  is, in reality, less than that of the actual  $\rho_+$ , giving a higher value of  $\tau_{pp}$ . The accurate value of  $\tau_{pp}$  can be calculated using the cosmic ray energy spectrum and coupling the relativistic and non-relativistic part of the accreting plasma which will be presented elsewhere. Nevertheless, our rough estimation assures that even with the accurate value of  $\tau_{pp}$  (which is higher than that of used here), the conditions of shock formation are highly satisfied.

If we now assume that a fraction  $\mathcal{E}_F$  of the infalling energy is converted into radiation through the hadronic collision ( $p - p$ ) and mesonic ( $\pi^\pm, \pi^0$ ) decay, this  $\mathcal{E}_F$  will allow convergent steady state solutions and in the case of quasi-spherical infall, allows the development and maintenance of a standing, collisionless shock at a fixed distance  $r_{sh}$  measured in the unit of Schwarzschild radius.

The luminosity obtained from this energy  $\mathcal{E}_F$  at the shock location  $r_{sh}$  is assumed to be  $\mathcal{L}_1$ . For a spherical shock surface and taking care of neutrino losses, the expression for  $\mathcal{L}_1$  can be written as (KE86)

$$\mathcal{L}_1 = 2\pi r_{sh}^2 \rho_{sh} m_p u_{sh}^3 \mathcal{E}_F \quad (7a)$$

where  $m_p$  is the mass of the proton and subscript  $sh$  indicates that the respective quantities are measured at the shock location.

If we assume that the pressure of the relativistic particles  $\mathcal{P}_{rel}$  (uniform inside the shock), the average energy density is then  $3\mathcal{P}_{rel}$ , so alternatively the luminosity can be expressed in terms of the volume integral of the emissivity  $\epsilon$  due to the hadronic ( $p - p$ ) collision where

$$\epsilon = \frac{3\mathcal{P}_{rel}}{\tau_{pp}}$$

Thus the alternative expression for the luminosity obtained as function of  $\mathcal{P}_{rel}$  would be,

$$\mathcal{L}_2 = \frac{4\pi r_{sh}^3 \mathcal{P}_{rel}}{\tau_{pp}} \quad (7b)$$

Defining  $\delta$  as the ratio of downstream relativistic particle pressure to incoming ram pressure at the shock, we obtain

$$\mathcal{P}_{rel} = \delta \rho_{sh} m_p u_{sh}^2 \quad (8)$$

Equating (7a) and (7b) and putting the values of  $\rho, \mathcal{P}_{rel}$ , and  $\tau_{pp}$  from eqs. (3), (6a) and (8) respectively, we get the expression for the shock location as a function of various inflow parameters as

$$r_{sh} = \frac{3\sigma_{pp} c \dot{M}_{in}}{u_{sh}^2 \Theta_{in}} \left( \frac{\delta}{\mathcal{E}_F} \right) \quad (9)$$

The ratio  $\left( \frac{\delta}{\mathcal{E}_F} \right)$  as a function of shock Mach number  $M_{sh}$  for a high shock Mach number solution (low energy inflow) is obtained from the empirical solution deduced by Ellison and Eichler (Ellison & Eichler, 1984), after suitable modification required for our model.

## 2.2 Outflow model

In ordinary stellar mass loss computations (Tarafer, 1988 and references therein) the outflow is assumed to be isothermal till the sonic point. This assumption is probably justified, since copious photons from the stellar atmosphere deposit momenta on the slowly outgoing and expanding outflow and possibly make the flow close to the isothermal. This need not be the case for outflow from black hole candidates. Our *effective boundary layer*, being pretty close to the black hole, are very hot and most of the photons emitted may be swallowed by the black hole itself instead of coming out of the region and depositing momentum onto the outflow.

Thus, the outflow could be cooler than the isothermal flow in our case. We choose polytropic outflow with a different polytropic index  $\gamma_o < \gamma$  due to momentum deposition. As a fraction of infalling energy density ( $\mathcal{E}_F$ ) is converted into radiation, specific energy of the outflow is somewhat less than that of the inflow. Nevertheless, the outflow specific energy is also kept constant throughout the flow.

The following two conservation laws are valid for the outflow :

$$\mathcal{E}' = \frac{v(r)^2}{2} + n' a(r)^2 - \frac{1}{2(r-1)} \quad (10)$$

$$\dot{M}_{out} = \Theta_{out} \rho(r) v(r) r^2 \quad (11)$$

Where  $\mathcal{E}'$  is the specific energy of the outflow and  $\mathcal{E}' < \mathcal{E}$  and  $n' = (\gamma_o - 1)^{-1}$  is the polytropic constant of the outflow.  $\Theta_{out}$  is the solid angle subtended by the outflow and  $v(r)$  is the velocity of the outflow.

For simplicity of calculation, we assume that the outflow is also quasi-spherical and  $\Theta_{out} \approx \Theta_{in}$ . Defining  $\dot{R}_{in}$  as the mass outflow rate, we obtain

$$\dot{R}_{in} = \frac{\dot{M}_{out}}{\dot{M}_{in}} \quad (12)$$

It is obvious from the above discussion that  $\dot{R}_{in}$  should have some complicated functional dependences on the following parameters

$$\dot{R}_{in} = \Psi(\mathcal{E}, \dot{M}_{in}, r_{sh}, R_{comp}, \gamma, \gamma_o) \quad (12a)$$

### 2.3 Procedure to solve the inflow and outflow equations simultaneously

Before we proceed into further detail, a general understanding of the transonic inflow outflow system in present case is essential to understand the basic scheme of the solution procedure. Let us consider the transonic accretion first. Infalling matter becomes supersonic after crossing a saddle type sonic point, location of which is determined by the inflow parameters such as the specific energy  $\mathcal{E}$ ,  $\dot{M}_{in}$  (in the unit of Eddington rate) and  $\gamma$  of the *inflow*. This supersonic flow then encounters a shock (if present) location of which ( $r_{sh}$ ) is determined from eqs. 9. At the shock, part of the incoming matter, having higher entropy density, is likely to return back as wind through a sonic point other than the one through which it just entered. Thus a combination of transonic topologies, one for the inflow and other for the outflow (passing through a different sonic point and following the topology *completely different* than that of the “self-wind” of the accretion), is required to obtain a full solution. So it turns out that finding a complete set of self-consistent inflow outflow solutions ultimately boils down to locating the sonic point of the *outflow* and the mass flux through it. Thus a supply of parameters  $\mathcal{E}$ ,  $\dot{M}_{in}$ ,  $\gamma$  and  $\gamma_o$  make a self consistent computation of  $\dot{R}_{in}$  possible. Here  $\gamma_o$  is supplied as free parameter because the self-consistent computation of  $\gamma_o$  directly using  $\mathcal{E}$ ,  $\dot{M}_{in}$  and  $\gamma$  has not been attempted in this work, instead we put a constrain that  $\gamma_o < \gamma$  always and for any value of  $\gamma$ . In reality,  $\gamma_o$  is directly related to the heating and cooling procedure of the outflow.

The following procedure is adopted to obtain a complete so-

lution:

From eqs. (1) & (2), we get the expression for the derivative

$$\frac{du(r)}{dr} = \frac{\frac{2a(r)^2}{r} - \frac{1}{2(r-1)^2}}{u(r) - \frac{a(r)^2}{u(r)}} \quad (13)$$

At the sonic point, the numerator and denominator separately vanish and give rise to the so called sonic point condition;

$$u_c = a_c = \frac{\sqrt{r_c}}{2(r_c - 1)} \quad (14)$$

Where the subscripts  $c$  represents the quantities at the sonic point. The derivative at the sonic point  $\left(\frac{du}{dr}\right)_c$  is computed using the L' Hospital rule. The expression for  $\left(\frac{du}{dr}\right)_c$  is obtained by solving the following polynomial,

$$\left(\frac{2n+1}{n}\right) \left(\frac{du}{dr}\right)_c^2 + \frac{3u_c}{nr_c} \left(\frac{du}{dr}\right)_c - \left\{ \frac{1}{(r_c-1)^3} - \frac{2a_c^2}{r_c} \left(\frac{n+1}{n}\right) \right\} = 0 \quad (15)$$

Using fourth order Runge Kutta method,  $u(r)$  and  $a(r)$  are computed along the inflow from the *inflow* sonic point till the position where the shock forms. The shock location is calculated by simultaneously solving the eqs. (1), (2) and (9). With the known value of  $\mathcal{E}'$  and  $\gamma_o$ , it is easy to compute the location of the sonic point of the *outflow* from eqs. (10) and (11). At the outflow sonic point, the outflow velocity  $v_c$  and polytropic sound velocity  $a_c$  is computed in the same manner performed for the inflow. Using eqs. (10) and (11),  $\left(\frac{dv}{dr}\right)_c$  and  $\left(\frac{du}{dr}\right)_c$  is computed as was done for the inflow. Runge-Kutta method is then employed to integrate from the *outflow* sonic point towards the black hole to find out the outflow velocity  $v$  and density  $\rho$  at the *shock location*. The outflow rate is then computed using eqs.(12).

## 3 RESULTS

Fig. 1 shows a typical solution which combines the accretion and the outflow. The input parameters are  $\mathcal{E} = 0.001$ ,  $\dot{M}_{in} = 1.0$  Eddington rate ( $\mathcal{E}_d$  stands for the Eddington rate in the figure) and  $\gamma = \frac{4}{3}$  corresponding to relativistic inflow. The solid curve with an arrow represents the pre-shock region of the inflow and the solid vertical line with double arrow at  $X_{pps}$  (the subscript *pps* stands for *pair plasma mediated shock*) represents the shock transition. Location of shock is obtained using the eqs. 4 for a particular set if inflow parameters mentioned above. Three dotted curves show the three different outflow branches corresponding to different polytropic index of the outflow as  $\gamma_o = 1.3$  (left most curve), 1.275 (middle curve) and 1.25 (rightmost curve). It is evident from the figure that the outflow moves along the solution curves completely different from that of the “wind solution” (solid line marked with an outward directed arrow) of the inflow which passes through the sonic point  $P_s$ . The mass loss ratio  $\dot{R}_{in}$  for these cases are 0.0023, 0.00065 and 0.00014 respectively. In Fig. 2., we have plotted the variation of  $\dot{R}_{in}$  with incoming specific energy  $\mathcal{E}$  for a set of values of  $\dot{M}_{in}$  (measured in the unit of Eddington rate shown as  $\mathcal{E}_d$  in

the Fig.) shown in the Fig. It is observed that  $R_{\dot{m}}$  monotonically increases with energy. This is because as  $\mathcal{E}$  increases keeping the Eddington rate of the inflow fixed, the shock Mach number  $M_{sh}$  decreases result of which is the decreament of shock location  $r_{sh}$  and post shock density (via eqs. 5.) but the increament of the post shock fluid velocity ( $v_{sh}$ ) with which the matter leaves the shock surface. The outflow rate  $R_{\dot{m}}$ , which is the product of these three quantities, in general increases monotonically with  $\mathcal{E}$  due to the combined tug of war of these three quantities. Moreover, closer the shock forms to the black hole, the greater will be the amount of gravitational potential available to be put onto the relativistic hadrons to provide more outward pressure at the shock boundary which gives a stronger “kick” to the accreting matter, the result of which is the increament in  $R_{\dot{m}}$ . All these points are manifested in Fig. 3 where we have shown the variation of  $R_{\dot{m}}$  as a function of compression ratio  $R_{comp}$  (solid curve), the shock location  $r_{sh}$  (dotted curve) and the injection velocity of the outflow  $v_{sh}$  (dashed curve). The fig. is drawn for a fixed  $\gamma = \frac{4}{3}$  and  $\gamma_o = 1.3$   $R_{comp}$  and  $v_{rsh}$  are scaled as  $R_{comp} \rightarrow (R_{comp} - 5.890) \times 10^3$  and  $v_{rsh} \rightarrow 4 \times 10^{-6} v_{rsh}$ . The unequal gaps between the curves with different  $\mathcal{E}_d$  implies that when inflow energy  $\mathcal{E}$  is kept fixed,  $R_{\dot{m}}$  nonlinearly increases with Eddington rate. This is because, as  $\mathcal{E}$  is kept fixed while  $\dot{M}_{in}$  is varied, the amount of infalling energy converted to produce high energy protons is also fixed, so higher is the value of  $\dot{M}_{in}$  (in the unit of Eddington rate), the larger is the distance of the shock surface (measured from the black hole) and the outflowing matter feels low inward gravitational pull, the result of which is the non-linear corelation of  $R_{\dot{m}}$  with  $\dot{M}_{in}$ .

To have better insight of the behavior of the outflow, in Fig. 4a. and 4b, we plot  $R_{\dot{m}}$  as a function of the polytropic index of the inflow  $\gamma$  (Fig. 4a) and that of the outflow  $\gamma_o$  (Fig. 4b) for fixed  $\mathcal{E} = 0.001$  and  $\dot{M}_{in} = 1.0\mathcal{E}_d$ . The range of  $\gamma$  shown here are the range for which shock forms for the specified  $\mathcal{E}$  and  $\dot{M}_{in}$ . In Fig. 4b, it is observed that shock forms for the values of  $\gamma_o$  lower than that shown in the figure, but mass loss rate is then so small that we did not plot for  $\gamma_o < 1.23$ . The general conclusion is that  $R_{\dot{m}}$  correlates with  $\gamma_o$ . This is because as  $\gamma_o$  increases, shock location and post shock density of matter does not change (as  $\gamma_o$  does not have any role in shock formation or in determining the  $R_{comp}$ ) but the sonic point of the outflow is pushed inward, hence the velocity with which outflow leaves the shock surface goes up resulting the increament in  $R_{\dot{m}}$ . However,  $R_{\dot{m}}$  anticorrelates with  $\gamma$  which is observed from Fig. 4a.

#### 4 CONCLUDING REMARKS

In the present paper, we have computed the mass outflow rate from the relativistic low energy matter quasi-spherically accreting onto Schwarzschild type black holes. The free parameters chosen for the inflow are the specific energy  $\mathcal{E}$ , mass inflow rate  $\dot{M}_{in}$  (In the unit of Eddington rate) and the polytropic index  $\gamma$  of the inflow. Only one extra parameter which was supplied for the outflow is its polytropic index  $\gamma_o$ . We have computed the mass outflow rate  $R_{\dot{m}}$  taking the help of inflow parameters only (except  $\gamma_o$ ) thus could analytically connect the accretion and wind type topologies

self-consistently. In our computation, we could investigate the dependence of  $R_{\dot{m}}$  as a function of different physical quantities governing the inflow.

The basic conclusions of this paper are the followings,

- (i) It is possible that outflows for quasi-spherical Bondi type accretion onto a Schwarzschild black hole are coming from the pair plasma pressure mediated shock surface.
- (ii) The outflow rate monotonically increases with the specific energy of the inflow and nonlinearly increases with the Eddington rate of the infalling matter.
- (iii)  $R_{\dot{m}}$ , in general, correlates with  $\gamma_o$  but anticorrelates with  $\gamma$ .
- (iv) Generally speaking, as our model deals with high shock Mach number (low energy accretion) solutions, outflows in our work always generate from the supersonic branch of the inflow, i.e, shock is always located *inside* the sonic point.
- (v) Unlike the mass outflow from the *accretion disks* around black holes (Das, 1998; 1998a; DC98) here we found that the value of  $R_{\dot{m}}$  is distinguishably small. This is because matter is ejected out due to the pressure of the relativistic plasma pairs which is *less enough* in comparison to the pressure generated due to the presence of significant angular momentum. However, in the present work we have dealt only high Mach number solution which means matter is accreting with very low energy (cold inflow, as it is described in literature). This is another possible reason to obtain a low mass loss rate. If, instead of high Mach number solution, we would use low Mach number solution, e.g, high energy accretion, the mass outflow would be considerably higher (this is obvious because it has already been established in present work that  $R_{\dot{m}}$  increases with  $\mathcal{E}$ . In our next work, we will present this type of model by calculating  $R_{comp}$  and  $(\frac{\delta}{\mathcal{E}_F})$  for low Mach number solution.

In the literature, we did not find any numerical simulation work which deal with the type of outflow we discuss in this paper (outflow from zero angular momentum inflow). On the other hand, observationally it is very difficult to calculate the outflow rate exactly from a real system, as it depends on too many uncertainties, such as filling factors and projection effects. So, at present, we don't have any offhand results in this field with which our result could be compared.

There are a number of possible improvements which could be made on this preliminary work. For instance, the effect of radiation pressure on both the inflow and the outflow could be taken into account. A preliminary investigation shows that the effect of radiation force, when included into the basic conservation equations (eqs. (1), (2) and (10), (11)) decreases the value of  $R_{\dot{m}}$ . This is probably because introduction of any radiation term which is proportional to  $\frac{1}{r^2}$ , weakens gravity and pushes the shock location outwards, combined effect of which is the decreament of  $R_{\dot{m}}$ . Another possible improvement is to include the magnetic field to give the outflow an appropriate geometry. In our model, we assumed the outflow to be quasi-spherical as like the inflow. The introduction of magnetic field probably could collimate the outflow thus would help to investigate the structure of jet related to the model of AGN without accretion disk in greater detail. Lastly, we did not self consistently compute  $\gamma_o$  as a function of inflow parameters and  $\gamma_o$  was supplied as

a free parameter. In our future work, we will be presenting the self consistent calculation of  $\gamma_o$  to reduce the number of free parameters of the problem.

So far, we made the computation around Schwarzschild black hole. Our work could be extended to study mass outflow in kerr space time using pseudo-Kerr potential (Chakrabarti & Khanna, 1992) and incorporating the frame dragging effect. This is under preparation and will be presented elsewhere.

### Acknowledgement :

The author expresses his gratitude to Prof. S. K. Chakrabarti and his colligues Mr. A. Ray for constructive discussions and helpfull suggestions which improved the quality of this work.

### REFERENCES

- Axford, W. I., Leer, E., Skadron, G., 1977, *Proc. 15th International Cosmic Ray Conf. (plovdiv)* **11** 132.
- Bondi, H., 1952, *Mon. Not. Roy Astron. Soc.* **112** 195.
- Chakrabarti, S. K., 1998, in *Observational Evidence for Black Holes in the Universe*, Ed. S. K. Chakrabarti (Kluwer Academic: Holland), **p.19** (also in *astro-ph/9807104*)
- Chakrabarti, S. K., Khanna, R., 1992, *Mon. Not. Roy Astron. Soc.* **256** 300.
- Das, T. K., 1998, in *Observational Evidence for Black Holes in the Universe*, Ed. S. K. Chakrabarti (Kluwer Academic: Holland), **p. 113** (also in *astro-ph/9807105*)
- Das, T. K., 1998a, *Proc-Xth International Summer School Seminar on Recent Problems in Theoretical & mathematical Physics. Kazan, Russia. June 18 - July 3 '98.*
- Das, T. K., 1998b, *Proc-International Mini Workshop on Mathematics and Mathematical Physics. Sept 1 - 3, '98.*
- Das, T. K., Chakrabarti, S. K., 1998, *Astrophys. J. submitted.* (*astro-ph/9809109*)
- Eichler, D., 1979, *Astrophys. J.* **232** 106.
- Ellision, D. C., Eichler, D., 1984, *Astrophys. J.* **286** 691.
- Ellision, D. C., Eichler, D., 1985, *Phys. Rev. Letters* **55** 2735.
- Flammang, R. A., 1982, *Mon. Not. Roy Astron. Soc.* **199** 833.
- Kazanas, D., Ellison, D. C., 1986, *Astrophys. J.* **304** 178.
- Meszaros, P., Ostriker, J. P., 1983, *Astrophys. J.* **273** L59.
- Paczyn'ski, B., Wiita, P. J., 1980, *Astn & Astph* **88** 23.
- Protheroe, R. J., Kazanas, D., 1983, *Astrophys. J.* **265** 620.
- Rees, M., 1977, *QJRAS*. **18** 429.
- Tarafdar, S. P., 1988, *Astrophys. J.* **331** 932.

### Figure Captions

Fig. 1: Solution topology for three different  $\gamma_o$  (1.3, 1.275, 1.25) for  $\mathcal{E} = 0.001$ ,  $\dot{M}_{in} = 1.0$ ,  $\mathcal{E}_d = 1.0$ ,  $\gamma = \frac{4}{3}$ .  $P_s$  indicates the sonic point of the *inflow* where  $X_{pps}$  stands for the shock location. See text for details.

Fig. 2: Variation of  $R_{in}$  with *inflow* specific energy  $\mathcal{E}$  for different Eddington rate (marked as  $\mathcal{E}_d$  in the figure). Other parameters are  $\gamma = \frac{4}{3}$  and  $\gamma_o=1.3$ .  $R_{in}$  monotonically increases with  $\mathcal{E}$ , see text for detail.

Fig. 3: Variation of  $R_{in}$  with the compression ratio  $R_{comp}$  (solid curve), shock location  $X_{ps}$  (dotted curve) and *outflow* velocity at shock  $v_{sh}$  (dashed curve). Other parameters are  $\mathcal{E}=0.001$ ,  $\dot{M}_{in}=1.0$ (In the unit of Eddington rate),  $\gamma = \frac{4}{3}$  and  $\gamma_o=1.3$ .

Fig. 4a: Variation of  $R_{in}$  with polytropic index of *inflow*  $\gamma$ . Other parameters are  $\mathcal{E}=0.001$ ,  $\dot{M}_{in}=1.0$  (in the unit of Eddington rate) and  $\gamma_o=1.3$ .

Fig. 4b: Variation of  $R_{in}$  with polytropic index of *outflow*  $\gamma_o$ . Other parameters are  $\mathcal{E}=0.001$ ,  $\dot{M}_{in}=1.0$  (in the unit of Eddington rate) and  $\gamma = \frac{4}{3}$ .











